

**The Bath Ankylosing Spondylitis Radiology Index** for the spine is applied to anteroposterior and lateral lumbar radiographs and lateral cervical radiographs. For the lumbar spine, defined as the lower border of T12 to the upper border of S1, anteroposterior and lateral radiographs are examined together and the view showing more significant change is scored. For the cervical spine, defined as the lower border of C1 to the upper border of C7, only the lateral view is scored.

Score	Grade	System applies to both the lumbar and the cervical spine (grade each as 0 – 4)
0	Normal	No change
1	Suspicious	No definite change
2	Mild	Any number of erosions, squaring, or sclerosis, with or without syndesmophytes, on less than or equal to 2 vertebrae
3	Moderate	Syndesmophytes on greater than or equal to 3 vertebrae, with or without fusion involving 2 vertebrae
4	Severe	Fusion involving greater than or equal to 3 vertebrae

#### **Scoring Instructions**

The Bath Ankylosing Spondylitis Radiology Index score is calculated by adding the average score for the sacroiliac joints to the scores for the lumbar and cervical spine.

Puntuación del BASRI aplicado a columna cervical/lumbar y caderas

Puntuación	Grado	Descripción
<i>Columna cervical/lumbar</i>		
0	Normal	Sin alteraciones
1	Sospechosa	Sin cambios definitivos
2	Leve	Alguna erosión, cuadratura o esclerosis con o sin sindesmofitos en 2 o menos vértebras
3	Moderada	Sindesmofitos en 3 o más vértebras con o sin fusión afectando a 2 vértebras
4	Severa	Fusión que afecta a 3 o más vértebras
<i>Caderas</i>		
0	Normal	Sin alteraciones
1	Sospechosa	Sin cambios definidos o estrechamiento focal del espacio articular
2	Leve	Estrechamiento concéntrico del espacio articular > 2 mm
3	Moderada	Estrechamiento concéntrico del espacio articular ≤ 2 mm o aposición hueso a hueso < 1 cm
4	Severa	Deformidad ósea o aposición hueso a hueso ≥ 1 cm