

Referencias bibliográficas

Belenguer R, Ramos-Casals M, Brito-Zerón P, del Pino J, Sentís J. Influence of clinical and immunological parameters on the health quality of life of patients with primary Sjögren's syndrome. Clinical and Experimental Rheumatology. 2005; 23: 351-356.

Bullo V et al. Nordic Walking can be incorporated in the exercise prescription to increase aerobic capacity, strength and quality of life for elderly: a systematic review and meta-analysis. Rejuvenation research. 2017; <https://doi.org/10.1089/rej.2017.1921>

Champey J, Corruble E, Gottenberg JE, Buhl C, Meyer T, Caudmont C et al. Quality of life and psychological status in patients with primary Sjögren's Syndrome and Sicca Symptoms without autoinmunes features. Arthritis & Rheumatism (Arthritis Care & Research). 2006; 55(3): 451-457.

Ejercicio físico, aliado de los pacientes con Síndrome de Sjögren, disponible en <https://www.efesalud.com/ejercicio-fisico-aliado-pacientes-sindrome-sjogren/> (consultado 27-8-2020)

González C, Mora M. Nordic Walking through cancer. An exercise prescription model to improve survivorship. Comunicación escrita en el European Cancer Survivorship & Rehabilitation Symposium. 2018. Copenhague (Dinamarca).

Hagner-Derengowska M et al. The effect a 10-week nordic walking training program on the level of GH and LH in elderly women. Climacteric. 2015; 18(6): 835-840.

Iveta N et al. Effects of Nordic Walking on cardiovascular performance and quality of life in coronary artery disease. European Journal of physical and rehabilitation medicine. 2020. DOI: 10.23736/S1973-9087.20.06120-1

Juarez M, Toms TE, de Pablo P, Mitchell S, Bowman S, Nightingale P et al. Cardiovascular risk factors in women with primary Sjögren's syndrome: United Kingdom primary Sjögren's syndrome registry results. Arthritis Care & Research. 2014; 66(5): 757-764.

Karageorgas T, Fragioudaki S, Nezos A, Karaiskos D, Moutsopoulos HM, Mavragani CP. Fatigue in primary Sjögren's Syndrome: clinical, laboratory, psychometric and biological associations. Arthritis Care & Research. 2016; 68(1): 123-131.

López-Valiente, C et al. Beneficios del ejercicio físico en pacientes con Síndrome de Sjögren primario: estudio de un caso. E-motion: Revista de Educación, Motricidad e Investigación. 2018; 10: 1-10.

Mengshoel AM, Norheim KB, Omdal R. Primary Sjögren's Syndrome: fatigue is an ever-present, fluctuating un uncontrollable lack of energy. *Arthritis Care & Research.* 2014; 66(8): 1227-1232.

Miyamoto ST et al. Supervised walking improves cardiorespiratory fitness, exercise tolerance, and fatigue in women with primary Sjögren's syndrome: a randomized-controlled trial. *Rheumatol Int.* 2019; 39(2): 227-238.

Nascimiento de Carvalho B, Torres dos Reis-Neto ET, Jennings F, Sardinha U, Kadratz F, Helder K et al. The relevance of performing exercise test before starting supervised physical exercise in asymptomatic cardiovascular patients with rheumatic diseases. *Rheumatology.* 2016; 55 (11): 1978-1986.

Nordic Walking y Síndrome de Sjögren, disponible en <https://4trebol.com/2020/05/26/nordic-walking-y-sindrome-de-sjogren/#more-9302> (consultado 28-8-2020)

Theander L, Strömbeck B, Mandl T, Theander E. Sleepiness or fatigue? Can we detect treatable causes of tiredness in primary Sjögren's syndrome?. *Rheumatology.* 2010; 49: 1177-1183.

Strömbeck BE, Theander E, Jacobsson TH. Effects of exercise on aerobic capacity and fatigue in women with primary Sjögren's syndrome. *Rheumatology.* 2007; 46: 868-87.

Szeffler -Derela J et al. Effectiveness of 6-week nordic walking training on functional performance, gait quality and quality of life in Parkinson's disease. *Medicina.* 2020; 56 (7): 356.

Vehí C et al. Nordic walking for cardiovascular prevention in patients with ischemic heart disease or metabolic syndrome. *Med Clin (Barcelona).* 2016; 147(12): 537-539.

Vílchez ME, Calvo-Arencibia A. Evidencia científica de la marcha nórdica en fisioterapia: revisión bibliográfica. *Fisioterapia.* 2016; 38(5): 251-264.

White PD, Goldsmith KA, Johnson AL, Potts L, Walwyn R, DeCesare JC et al. Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial. *Lancet.* 2011; 377(9768): 823-836.