

**Responsiveness in Rheumatoid Arthritis. A Report from the OMERACT 11
Ultrasound Workshop.**

Iagnocco A1, Naredo E, Wakefield R, Bruyn GA, Collado P, Jousse-Joulin S, Finzel S, Ohrndorf S, Delle Sedie A, Backhaus M, Berner-Hammer H, Gandjbakhch F, Kaeley G, Loeuille D, Moller I, Terslev L, Aegerter P, Aydin S, Balint PV, Filippucci E, Mandl P, Pineda C, Roth J, Magni-Manzoni S, Tzaribachev N, Schmidt WA, Conaghan PG, D'Agostino MA. J Rheumatol 2014 Feb; 41(2):379-82

**Grupo de Trabajo de Ecografía de la Sociedad Española de Reumatología
(ECOSER). Ultrasound in rheumatology: Where are we and where are we going
De Miguel E, Andreu JL, Naredo E, Möller I, Grupo de Trabajo de Ecografía de la
Sociedad Española de Reumatología (ECOSER).**

Reumatol Clin. Reumatol Clin. 2014;10(1):6-9 Financiado

**Cardiovascular risk stratification in rheumatic diseases: carotid ultrasound is
more sensitive than Coronary Artery Calcification Score to detect subclinical
atherosclerosis in patients with rheumatoid arthritis.**

Corrales A1, Parra JA, González-Juanatey C, Rueda-Gotor J, Blanco R, Llorca J, González-Gay MA. Ann Rheum Dis. 2013; 72(11):1764-70. Financiado

**Carotid ultrasound is useful for the cardiovascular risk stratification of patients
with rheumatoid arthritis: results of a population-based study.**

Corrales A1, González-Juanatey C, Peiró ME, Blanco R, Llorca J, González-Gay MA
Ann Rheum Dis. 2013; Financiado

**Current state of musculoskeletal ultrasound in paediatric rheumatology: results of
an international survey**

Magni-Manzoni S1, Collado P, Jousse-Joulin S, Naredo E, D'Agostino MA, Muratore V, Merli P, Roth J; on behalf of the Paediatric Ultrasound Group of the OMERACT Ultrasound Task Force. Rheumatology (Oxford). 2013;

**Dynamic ultrasound assessment of medial meniscal subluxation in knee
osteoarthritis**

Acebes C, Romero FI, Contreras MA, Mahillo I, Herrero-Beaumont G. Rheumatology (Oxford). 2013; 52(8):1443-7

**Patient self-assessment and physician's assessment of rheumatoid arthritis
activity: which is more realistic in remission status? A comparison with
ultrasonography.**

Rheumatology (Oxford). 2013;52(12):2243-50.

**Reduced joint assessment vs comprehensive assessment for ultrasound detection of
synovitis in juvenile idiopathic arthritis**

Collado P, Naredo E, Calvo C, Gamir ML, Calvo I, García ML, et al. *Rheumatology (Oxford)*. 2013;52(8):1477-84

Three-dimensional volumetric ultrasound: a valid method for blinded assessment of response to therapy in rheumatoid arthritis.

Naredo E1, Acebes C, Brito E, de Agustín JJ, de Miguel E, Mayordomo L, Möller I, Moragues C, Rejón E, Rodríguez A, Uson J, Garrido J, Martínez-Hernández D; Ultrasound School of the Spanish Society of Rheumatology. *Rheumatol*. 2013;40(3):253-60

Ultrasound joint inflammation in rheumatoid arthritis in clinical remission: how many and which joints should be assessed?

Naredo E, Valor L, De la Torre I, Martínez-Barrio J, Hinojosa M, Aramburu F, Ovalles-Bonilla JG, Hernández D, Montoro M, González CM, López-Longo J, Monteagudo I, Carreño L. *Arthritis Care Res (Hoboken)*. 2013;65(4):512-7.

Ultrasound-detected musculoskeletal urate crystal deposition: which joints and what findings should be assessed for diagnosing gout?

Naredo E1, Uson J, Jiménez-Palop M, Martínez A, Vicente E, Brito E, Rodríguez A, Cornejo FJ, Castañeda S, Martínez MJ, Sanz J, Möller I, Batlle-Gualda E, Garrido J, Pascual E. *Ann Rheum Dis*. 2013; PMID: 23709244 No financiado